| Paediatric clinician trainee number | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|--|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| Basic Life Support/SVT | 46% | 54% | 38% | 61% | 36% | 41% | 38% | 41% | 52% | 61% | 61% | 55% |
| Neonatal Medicine | 81% | 92% | 70% | 89% | 81% | 65% | 73% | 65% | 76% | 96% | 84% | 89% |
| Acute Abdomen & Other surgical cases | 74% | 66% | 74% | 90% | 78% | 70% | 70% | 62% | 74% | 88% | 76% | 94% |
| Measles, VHF, Enteric Fever | 75% | 71% | 57% | 81% | 64% | 62% | 64% | 60% | 69% | 74% | 76% | 62% |
| Other Common Childhood Infections | 90% | 80% | 47% | 73% | 50% | 67% | 80% | 67% | 77% | 77% | 73% | 80% |
| Neck Lumps | 65% | 65% | 38% | 55% | 70% | 53% | 53% | 28% | 60% | 58% | 55% | 70% |
| Sickle Cell, Haemolytic Anaemia, Blood Clotting | 78% | 76% | 63% | 78% | 80% | 76% | 76% | 80% | 76% | 80% | 83% | 78% |
| Seizures | 88% | 92% | 74% | 84% | 90% | 74% | 82% | 72% | 80% | 86% | 76% | 90% |
| Acute Hepatitis & Liver Failure | 93% | 77% | 70% | 90% | 83% | 40% | 57% | 53% | 83% | 87% | 70% | 93% |
| GI Haemorrhage and Upper GI Disorders | 78% | 85% | 67% | 89% | 81% | 74% | 66% | 70% | 63% | 89% | 72% | 85% |
| Cancer in Children | 100% | 87% | 87% | 92% | 80% | 78% | 70% | 83% | 90% | 90% | 83% | 100% |
| Chronic Liver Failure/Vitamin & Mineral Deficiencies | 97% | 87% | 85% | 92% | 82% | 72% | 82% | 72% | 90% | 97% | 79% | 92% |
| Headaches | 95% | 95% | 80% | 98% | 90% | 88% | 90% | 80% | 90% | 95% | 88% | 93% |
| Persistent Diarrhoea, Malabsorption & Constipation | 95% | 88% | 84% | / | 86% | 81% | 79% | 81% | 81% | 91% | 86% | 93% |
| Neuromuscular Disorders & Spinal Cord Injury | 94% | 92% | 60% | 77% | 77% | 69% | 77% | 77% | 73% | 83% | 84% | 79% |
| Endocrine | 68% | 74% | 57% | 68% | 66% | 49% | 62% | 51% | 68% | 81% | 64% | 62% |